arXiv:1504.07826v3 [math.AP] 22 Sep 2015

Asymptotic stability of solitons to 1D Nonlinear Schrödinger Equations in subcritical case

Ze Li

u19900126@163.com

Wu Wen-Tsun Key Laboratory of Mathematics, Chinese Academy of Sciences
and
Department of Mathematics, University of Science and Technology of China

Abstract

In this paper, we prove the asymptotic stability of solitons to 1D nonlinear Schrödiger equations in subcritical case with some spectral assumptions. Although pure power nonlinearities are not included in our theorem because of the failure of spectral hypothesis, our work throws light on how to deal with weak nonlinearity. Since dispersive methods mostly need a high power nonlinearity, we develop a commutator method which dates back to the work of S. Cuccagna, V. Georgiev, N. Visciglia [4].

Keywords: nonlinear Schrödinger equations; asymptotic stability; subcritical case

1 Introduction

In this paper, we consider the following nonlinear schrödinger equation (NLS),

$$\begin{cases} iu_t = -\Delta u - F(|u|^2)u \\ u(h,x) = u_h(x). \end{cases} \tag{1.1}$$

where $u : [h,\infty) \times \mathbb{R} \to \mathbb{C}$. The asymptotic stability of solitons states that for any solution of NLS with initial data near some soliton tends to a modulated soliton and a dispersive part with a remainder small in time. This is best known for completely integrable equations for instance the cubic NLS in one dimension. The basic tool for this case is the inverse scattering. For general nonlinearities, the first asymptotic stability was obtained by A. Soffer and M. I. Weinstein [17] in context of the equation

$$iu_t = -\Delta u + [V(x) - |u|^{m-1}]u. \tag{1.2}$$

There has been a lot of succeeding works in this direction such as E. Kirr, A. Zarnescu [11], S. Gustafson, K. Nakanishi, T.P. Tsai [8], and M. Tetsu [18]. Since dispersive method mostly fails for subcritical nonlinearities, the asymptotic stability for (1.2) with subcritical nonlinearities is largely open. In this direction, E. Kirr, O. Mizrak [9] partly solves the problem in three dimensions for equation (1.2). Moreover, their results exclude the case $V = 0$.

In Buslaev and Perelman [3], under some spectral assumptions, they proved asymptotic stability for (1.1) with some special nonlinearities, indeed they require F to be a polynomial with degree more than four. Their work was extended to high dimensions in Cuccagna [5]. Then Perelman [13] and I. Rodnianski, W. Schlag, A. Soffer [14] established the asymptotic stability for multi-solitons in high dimensions was proved.

In another direction, asymptotic behaviors of solutions to super-mass critical NLS equations were studied. Since in this case, blow-up phenomenons occur, we can not expect asymptotic stability for all data. In [16], Schlag proved that locally around each ground-state soliton there exists a codimension-1 Lipschitz submanifold of $H^1(\mathbb{R}^3) \cap W^{1,1}(\mathbb{R}^3)$ of initial data that lead to global $H^1 \cap W^{1,1}$ solutions to NLS. These solutions tend to a modulated soliton and a dispersive term. Cuccagna [6] obtained a set of asymptotic stable solutions forming a codimension-1 initial data set without manifold structure for mass-super critical one dimensional NLS. Beceanu [1] studied cubic NLS in three dimensions, it was proved that there exists a codimension-1 real analytic manifold $\mathcal{N}$ such that for any initial data in $\mathcal{N}$ (1.1) has a forward global solution which decomposes into a moving soliton and a dispersive term.

There are three main factors affecting the study of asymptotic stability, namely potentials, dimensions and nonlinearities. Since the solitons of equation (1.2) arise as a perturbation of the eigenfunctions of the operator $-\Delta + V$, in contrast to the case of NLS equation, they have a fixed center, which makes the analysis partially easier. Besides the effects of the potential term, dimension plays an important role in proving asymptotic stability. Generally, because of the weaker decay of linearized operator in one dimension compared with high dimensions, the one dimension case is more difficult. The lower the power of nonlinearity is, the less decay we can gain, which makes the subcritical nonlinearity harder.

The degree restriction in [3] is used to offset the weak decay estimate of the linearized operator. Moreover, it seems impossible to remove the degree assumption only by dispersive estimates. In this paper, we aim to extend the result of [3] to subcritical nonlinearities. In order to deal with low degree nonlinearity, we develop a commutator method, which dates back to the work of S. Cuccagna, V. Georgiev, N. Visciglia [4].

First of all, we give the definition of solitons.

Definition 1.1. *When nonnegative radial function φ and $\sigma(t) = (\beta, \omega, b, v)$ satisfy*

$$\begin{aligned} &\Delta\varphi = \alpha^2\varphi/4 - |\varphi|^{p-1}\varphi \\ &\beta' = \omega, \omega' = 0, b' = v, v' = 0, \end{aligned} \tag{1.3}$$

then $w(x;\sigma) = exp(-i\beta + i\frac{1}{2}vx)\varphi(x - b; \alpha)$ is a solution to (1.1), where $\omega = \frac{1}{4}(v^2 - \alpha^2)$. We call $w(x;\sigma)$ solitons.

It is known that when $F(x) = |x|^{\frac{p-1}{2}}$, p is mass-subcritical, φ exists and it decays exponentially at infinity. (see Berestycki and Lions [2] where a larger class of nonlinearity was considered)

As in [3]. the linearization of (1.1) around the soliton $w(x, t; \sigma)$ is

$$i\partial_t\chi = -\Delta\chi + F(|w|^2)\chi + F'(|w|^2)w(w\chi + w\overline{\chi})$$

If we denote

$$\chi(x, t) = exp(i\Phi)f(y, t),\ \Phi = -\beta(t) + \frac{1}{2}vx,\ y = x - b,$$

then function f satisfies the equation

$$i\partial_t f = L(\alpha)f,$$

where

$$L(\alpha)f = -\Delta f + \alpha^2 f/4 + F(\varphi^2)f + F'(\varphi^2)\varphi^2(f + \overline{f}), \varphi = \varphi(y, \alpha). \tag{1.4}$$

For (1.4), we consider its complexification:

$$\begin{aligned} &i\partial_t\vec{f} = H(\alpha)\vec{f}, \vec{f} = (f, \overline{f})^t, \\ &H(\alpha) = H_0(\alpha) + V(\alpha), H_0(\alpha) = (-\Delta_y + \alpha^2/4)\theta_3, \\ &V(\alpha) = [F(\varphi^2) + F'(\varphi^2)\varphi^2]\theta_3 + iF'(\varphi^2)\varphi^2\theta_2, \end{aligned}$$

where θ_2 and θ_3 are the matrices:

$$\theta_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix},\ \theta_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

There are four known generalized eigenfunctions denoted by $\{\xi_1, \xi_2, \xi_3, \xi_4\}$ for eigenvalue zero to $H(\alpha)$, where the two radial functions are

$$\xi_1 = \begin{pmatrix} v_1 \\ \bar{v}_1 \end{pmatrix}, \xi_2 = \begin{pmatrix} v_2 \\ \bar{v}_2 \end{pmatrix}, v_1 = -i\varphi(y;\alpha), v_2 = -\frac{2}{\alpha}\varphi_\alpha(y;\alpha).$$

Moreover, $H(\alpha)\xi_1 = 0$, $H(\alpha)\xi_2 = i\xi_1$, $\langle \xi_1, \xi_2 \rangle = 0$.

We have the following assumptions about the nonlinearity F.

Assumption A:
(i) There exists a soliton with exponential decay. Equation (1.1) is globally wellposed in H^2 and the solution $u(t)$ satisfies $\|u(t)\|_{H^2} \le C(\|u_h\|_{H^2})$;
(ii) There exist $m, n > \frac{26}{7}$ such that for $k \in \{0,1,2,3,4\}$,

$$\left|\frac{d^k}{dx^k}F(x)\right| \le |x|^{\frac{m-1}{2}-k}, \text{ if } |x| \le 1,$$
$$\left|\frac{d^k}{dx^k}F(x)\right| \le |x|^{\frac{n-1}{2}-k}, \text{ if } |x| \ge 1.$$

(iii) The linearized operator $H(\alpha)$ has zero as its only eigenvalue with generalized eigenfunction space spanned by $\{\xi_1, \xi_2, \xi_3, \xi_4\}$, no resonance and no embedded eigenvalues in its continuous spectrum.

Remark 1.1 If we restrict ourself to power functions, roughly speaking, (ii) means the lowest degree of F is $\frac{19}{14}$, which improves the result in [3] where the degree of F is assumed to be at least four.

Remark 1.2 Although (iii) is widely assumed in the papers studying asymptotic stability, as we know, for pure power nonlinearity namely $F(x) = x^l$, (iii) fails. However there are some possible remedies to deal with the case when more than one discrete spectrum occurs, such as the so called " Fermi Golden Rule" hypothesis (FGR) introduced by Sigal [15].

Remark 1.3 If we assume $n < 5$, then (i) is naturally satisfied. Meanwhile (i) may holds for combined nonlinearities, hence we use (i) instead of $n < 5$.

Before going to the main theorem, we give a proposition on the wellposedness. The proof is presented in Appendix A.

Proposition 1.1. *For F satisfying Assumption A, if the initial data $u_h \in H^2$ satisfies $\|u_h(1+|x|)^2\|_2 < \infty$, then there exists a unique solution $u(t)$ to equation (1.1), furthermore, we have*

$$\|u(t)\|_{H^2} \le C, \ \|u(t)(1+|x|)\|_2 \le C(1+t), \ \|u(t)(1+|x|^2)\|_2 \le C(1+t^4).$$

Define $\|u\|_\Sigma = \|u\|_{H^2} + \||x|\,\partial_x u\|_2 + \left\||x|^2 u\right\|_2$. From now on, we only consider radial functions. Our main theorem is the following:

Theorem 1.2. *$w(x;\sigma_0(t))$ is a soliton solution to (1.1), $\sigma_0 = (\beta_0, \omega_0, 0, 0)$, satisfying $\frac{d}{d\alpha}\|\varphi\|_2^2 \neq 0$ at α_0, where $-\frac{1}{4}\alpha_0^2 = \omega_0$ (Hypothesis B). Suppose that h is sufficiently large. If the radial initial data u_h in Proposition 1.1 satisfies $\|u_h - w(x;\sigma_0(0))\|_\Sigma \ll 1$, then there exists a soliton $w(x;\sigma_+(t))$ such that $u(x,t)$ decomposes as*

$$u(x,t) = w(x;\sigma_+(t)) + \chi(t),$$

where $\|\chi(t)\|_\infty \leq Ct^{-s+1}$ with $s = \frac{7}{4}^+$.

The whole proof is divided into five steps. In the first step, we split the solution into a modulated soliton and a remainder term. Second, we introduce the commutator operator and impose some suitable orthogonal conditions to the remainder. Third, we give the precise definition of the fractional power of the linearized operator and construct the following inequality

$$\|\chi\|_\infty \leq Ct^{-s}\||J_V(t)|^s\chi\|_{L_t^\infty L_x^2}, \tag{1.5}$$

where $|J_V(t)|^s$ is the commutator operator having the form $U(t)t^sH(\alpha(t))^{s/2}U(-t)$ defined in Section 3. In step four, we derive the equation for $|J_V(t)|^s u$. In the last step, we apply the Strichartz estimates and bound $\||J_V(t)|^s\chi\|_{L_t^\infty L_x^2}$, which combined with inequality (1.5) imply the decay of the remainder term, thus accomplishing the proof of the main theorem.

The core of our proof is (1.5), because (1.5) fills the gap between the weak decay resulting from dispersive effects in one dimension and the desired decay to the remainder. In fact, if we use dispersive estimate to bound the remainder, the best rate is $t^{-\frac{1}{2}}$. But it even cannot provide the existence of the limit to the modulated soliton. However, (1.5) can supply a decay as fast as you need, which is the ultimate advantage compared with dispersive methods.

There are two main difficulties in the establishment of (1.5). The first is how to give a proper definition of $H(\alpha)^{s/2}$. We notice that it is not a trivial thing, since $H(\alpha)$ is not an m-accertive operator, and more importantly, zero is its eigenvalue. The second difficulty is how to prove (1.5). A similar estimate for Schrödinger operator is given in [4] by inverse scattering theory. There are two key differences in our case. First, in [4], no discrete spectrum occurs, however, $H(\alpha)$ enjoys four dimensional generalized eigenfunction space. Second, $H(\alpha)$ corresponds to the inhomogeneous Sobolev space. Although the last difference leads to a faster decay, the first distinction makes the problem more difficult.

To overcome aforementioned difficulties, we develop some functional analysis method, which highlights the power of functional calculus. Specifically speaking, in order to get (1.5), we abandon inverse scattering theory, instead, we translate the problem to the corresponding estimate of the inverse of $H(\alpha)^{s/2}$, which reduces the problem to resolvent estimates. As to resolvent estimates, we apply a perturbation technique to obtain the decay of resolvent $(\lambda - H(\alpha))^{-1}$, when λ is sufficiently large. Although we cannot prove a similar decay for small λ, we can still solve our problem since 0 is not a singular point in our definition of $H(\alpha)^{s/2}$. It is important that (1.5) remains valid only for χ satisfying $U(-t)\chi \in P_c(H(\alpha))$, where P_c is the projection to the continuous part of $H(\alpha)$. Hence orthogonal conditions should be added to χ. This condition is quite different from the usual, which is assumed to guarantee that the remainder belongs to the continuous spectral part of $H(\alpha)$. As soon as we establish (1.5), the main theorem follows from the study of modulation equation and Strichartz estimates.

The paper is organized as follows. In Section 2, we give the orthogonal conditions and derive the modulation equation. In Section 3, we define the fractional order derivative of the linearized operator and commutator operator $|J_V|^s$. In Section four, we get the equation of $|J_V|^s u$ and establish the nonlinear estimates from which the proof of Theorem 1.1 follows.

For convenience, $(f, \bar{f})^t$ is simplified as $\vec{f}$ or f.

2 Orthogonal conditions

Define

$$|J_V(t)|^s f = \begin{pmatrix} M(t) & \\ & M(-t) \end{pmatrix} \left(t^2 H(\alpha(t))\right)^{s/2} \begin{pmatrix} M(-t) & \\ & M(t) \end{pmatrix} f,$$

where $M(t) = e^{i|x|^2/4t}$. The definition of $(H(\alpha(t))^{s/2}$ is given in Section 3. Here, we only need the fact that $H(\alpha)^{s/2}\xi_i = 0$, and $(H^*(\alpha))^{s/2}\theta_3 = \theta_3 H(\alpha)^{s/2}$.

We assume the solution to (1.1) is of the following form:

$$u(x,t) = w(x;\sigma(t)) + \chi(x,t), w(x;\ \sigma(t)) = \exp(-i\beta(t))\varphi(x,\alpha(t)),$$

where $\sigma(t) = (\beta(t), \omega(t), 0, 0)$ is not a solution of (1.3) in general. Define $\chi(x,t) = exp(-i\beta(t))f(x,t)$, $\vec{f} = (f, \bar{f})^t$ and

$$U(t) = \begin{pmatrix} M(t) & \\ & M(-t) \end{pmatrix},$$

then we impose four orthogonal conditions to f, namely for $i = 1, 2$,

$$\left\langle U(-t)\vec{f}, \theta_3\xi_i \right\rangle = 0 \tag{2.6}$$

and

$$\left\langle |J_V|^s\vec{f}, U(t)\theta_3\xi_i \right\rangle = 0, \tag{2.7}$$

where ξ_i means $\xi_i(\alpha(t))$.

In fact, (2.7) is a consequence of (2.6). Since $(H^*(\alpha))^{s/2}\theta_3 = \theta_3 H(\alpha)^{s/2}$, it holds

$$\begin{aligned}\left\langle |J_V|^s \vec{f}, U(t)\theta_3\xi_i \right\rangle &= t^s \left\langle U(-t)\vec{f}, (H^*(\alpha(t)))^{s/2}\theta_3\xi_i \right\rangle \\ &= t^s \left\langle U(-t)\vec{f}, \theta_3 (H(\alpha(t)))^{s/2}\xi_i \right\rangle.\end{aligned}$$

From $H(\alpha)^{s/2}\xi_i = 0$, we obtain

$$\left\langle U(-t)\vec{f}, \theta_3 (H(\alpha))^{s/2}\xi_i \right\rangle = 0;$$

Thus the orthogonal conditions are reduced to

$$\left\langle U(-t)\vec{f}, \theta_3\xi_i \right\rangle = 0. \tag{2.8}$$

The existence of $\sigma(t)$ follows from the following lemma.

Lemma 2.1. *If h is sufficiently large, $\chi(h)$ is small in L^2 norm, then there exists $\sigma(t) = (\beta(t), \omega(t), 0, 0)$ such that (2.8) holds.*

Proof We first prove it for $t = h$, namely finding appropriate α and β such that

$$\left\langle \begin{pmatrix} u(x,h) - e^{-i\beta}\varphi(x;\alpha) \\ \overline{u(x,h)} - e^{i\beta}\varphi(x;\alpha) \end{pmatrix}, \begin{pmatrix} e^{-i\beta} & \\ & e^{i\beta} \end{pmatrix} U(h)\theta_3\xi_i(\alpha) \right\rangle = 0.$$

This solvability is a consequence of the non-singularity of the corresponding Jacobian. Indeed, using $|M(t) - 1| \leq C\frac{|x|^2}{t}$, χ_h is small in L^2, h is sufficiently large, the main term of the Jacobian is

$$\begin{pmatrix} e & 0 \\ 0 & s \end{pmatrix},$$

where $e = -i\frac{d}{d\alpha}\left\|\varphi(x;\alpha)\right\|_2^2$, $s = -\frac{2}{\alpha}i\frac{d}{d\alpha}\left\|\varphi(x;\alpha)\right\|_2^2$. At α_0, e and s are nonzero by Hypothesis B, thus we have proved our lemma when $t = h$.

Second we show the existence of $\sigma(t)$, but this follows from the same argument in the remark after Proposition 1.3.1 in [3].

2.1 The system of equations for parameters

Define $\beta(t) = \int_0^t \omega(\lambda)d\lambda + \gamma$, then we rewrite (1.1) in terms of f:

$$if_t = L(\alpha(t))f + N(\varphi, f) - \gamma' f - \gamma'\varphi + i\omega'\frac{2}{\alpha}\varphi_\alpha,$$

where $N(\varphi, f) = F(|\varphi + f|^2)(\varphi + f) - F(\varphi^2)\varphi - F(\varphi^2)f - F'(\varphi^2)\varphi^2(f + \bar{f})$. By (2.8), we have

$$\operatorname{Im}\langle f, M(t)v_i\rangle = 0. \tag{2.9}$$

Lemma 2.2. *If $\|f\|_\infty$ is sufficiently small. then*

$$|\gamma'| + |\omega'| \le C(\|f\|_\infty^2 + \|f\|_\infty^{m-1}\|f\|_m + \|f\|_\infty^{n-1}\|f\|_n + \frac{1}{t^2}\|f\|_\infty + \frac{1}{t}\|f\|_\infty).$$

Proof Differentiating (2.9), we get

$$\begin{aligned}&\operatorname{Im}\left\langle i\gamma'\varphi + \omega'\frac{2}{\alpha}\varphi_\alpha, M(t)v_i\right\rangle + \operatorname{Im}\langle f, M(t)\partial_\alpha v_i\rangle\,\omega'\frac{2}{\alpha} + \operatorname{Im}\langle -iL(\alpha(t))f, M(t)v_i\rangle \\ &+ \operatorname{Im}\langle -iN(\varphi, f), M(t)v_i\rangle + \operatorname{Im}\langle i\gamma' f, M(t)v_i\rangle + \operatorname{Im}\left\langle f, \frac{d}{dt}M(t)v_i\right\rangle = 0.\end{aligned} \tag{2.10}$$

Direct calculations give

$$\left|\left\langle f, \frac{d}{dt}M(t)v_i\right\rangle\right| \le t^{-2}\|f\|_\infty. \tag{2.11}$$

Because of the orthogonal conditions (2.8), equation $H(\alpha)\xi_2 = i\xi_1$ and the obvious commutator inequality, we have

$$\begin{aligned}|\operatorname{Im}\langle -iL(\alpha(t))f, M(t)v_i\rangle| &= \left|\left\langle H(\alpha(t))\vec{f}, U(t)\theta_3 v_i\right\rangle\right| \\ &= \left|\left\langle \vec{f}, U(t)H^*(\alpha(t))\theta_3 v_i\right\rangle\right| + \left|\left\langle \vec{f}, [U(t), H^*(\alpha(t))]\theta_3 v_i\right\rangle\right| \\ &= \left|\left\langle \vec{f}, U(t)\theta_3 H(\alpha(t))v_i\right\rangle\right| + \left|\left\langle \vec{f}, [U(t), H^*(\alpha(t))]\theta_3 v_i\right\rangle\right| \\ &= \left|\left\langle \vec{f}, [U(t), H^*(\alpha(t))]\theta_3 v_i\right\rangle\right| \le t^{-1}\|f\|_\infty.\end{aligned}$$

Combined with (2.10) and (2.11), it follows,

$$\left\|\begin{pmatrix} 0 & n \\ -n & 0 \end{pmatrix}\begin{pmatrix} \gamma' \\ \omega' \end{pmatrix}\right\| \le O_1(\|f\|_\infty)\left\|\begin{pmatrix} \gamma' \\ \omega' \end{pmatrix}\right\| + O_2(f) + t^{-1}\|f\|_\infty + t^{-2}\|f\|_\infty, \tag{2.12}$$

where O_1 is the linear term of f, O_2 is the quadratic term of f, and $O_2 \le C(|f|^m + |\varphi|^{m-2}|f|^2 + |f|^n + |\varphi|^{n-2}|f|^2)$. Thus we have proved our lemma.

2.2 Reduction to time-independent linearized operator

Fix time $t_1 > h$. Suppose that $\omega(t_1) = \omega_1$, $\beta(t_1) = \beta_1$. Define $\omega_1 = -\frac{1}{4}\alpha_1^2$, $\gamma_1 = \beta_1 - \omega_1 t_1$, $\Phi_1 = -\omega_1 t - \gamma_1$. Let $\chi = e^{i\Phi_1} g(x,t)$, then g satisfies

$$ig_t = L(\alpha_1)g + D.$$

The function D is given by

$$\begin{aligned} D &= D_0 + D_1 + D_2 + D_3 + D_4, \\ D_0 &= e^{-i\Omega}[-\gamma'\varphi(\alpha) + \frac{2i}{\alpha}\omega'\varphi_\alpha], \quad \Omega = \Phi_1 - \Phi, \\ D_1 &= [F(\varphi^2(\alpha)) + F'(\varphi^2(\alpha))\varphi^2(\alpha) - F(\varphi^2(\alpha_1)) - F'(\varphi^2(\alpha_1))\varphi^2(\alpha_1)]g, \\ D_2 &= F'(\varphi^2(\alpha))\varphi^2(\alpha)[e^{-2i\Omega} - 1]\bar{g}, \\ D_3 &= [F'(\varphi^2(\alpha))\varphi^2(\alpha) - F'(\varphi^2(\alpha_1))\varphi^2(\alpha_1)]\bar{g}, \\ D_4 &= e^{-i\Omega}N(\varphi(\alpha), e^{i\Omega}g). \end{aligned}$$

The equation for $\vec{g} = (g, \bar{g})^t$ is

$$i\vec{g}_t = H(\alpha_1)\vec{g} + \vec{D}. \tag{2.13}$$

3 Fractional order derivative of $H(\alpha)$

For $\alpha > 0$, recall the linearized operator $H(\alpha)$. Define $\tau = \frac{1}{4}\alpha^2$ and

$$H_0 = \begin{pmatrix} -\Delta + \tau & \\ & \Delta - \tau \end{pmatrix}.$$

With these notations, the potential term in $H(\alpha)$ is given by $H(\alpha) = H_0 + V$, where V can be written as

$$\begin{pmatrix} V_1 & -V_2 \\ V_2 & -V_1 \end{pmatrix}.$$

The explicit formula for the kernel of H_0 and Young's inequality yield

Lemma 3.1. *For* $1 \le p \le \infty$, $\lambda \in \mathbb{C}\backslash[(-\infty, -\tau]\bigcup[\tau, \infty)]$,

$$\left\|(H_0 - \lambda)^{-1}\right\|_{p\to p} \le c_1[(\mathrm{Re}\sqrt{-\lambda + \tau})^{-2} + (\mathrm{Re}\sqrt{\lambda + \tau})^{-2}];$$

and for $2 \le p \le \infty$*,*

$$\Big\|(H_0-\lambda)^{-1}\Big\|_{2\to p} \le c_1[(\mathrm{Re}\sqrt{\tau-\lambda})^{-\frac{3}{2}-\frac{1}{p}} + (\mathrm{Re}\sqrt{\lambda+\tau})^{-\frac{3}{2}-\frac{1}{p}}].$$

Lemma 3.2. *Let* $\lambda \in \mathbb{C}\backslash[(-\infty,-\tau]\bigcup[\tau,\infty)]$*. If* $[\mathrm{Re}\sqrt{(-\lambda+\tau)}]^2 \ge 4c_1\|V\|_\infty$ *and* $[\mathrm{Re}\sqrt{(\lambda+\tau)}]^2 \ge 4c_1\|V\|_\infty$*,* $1 \le p \le \infty$*, then*

$$\|(H(\alpha)-\lambda)^{-1}\|_{p\to p} \le c[(\mathrm{Re}\sqrt{\tau-\lambda})^{-2} + (\mathrm{Re}\sqrt{\lambda+\tau})^{-2}],$$

where c *is independent of* V*.*

Proof Formally we have

$$(H(\alpha)-\lambda)^{-1} = (I+(H_0-\lambda)^{-1}V)^{-1}(H_0-\lambda)^{-1}.$$

Precisely, if $\Big\|(H_0-\lambda)^{-1}V\Big\|_{p\to p} < \frac{1}{2}$, then by Neumann series the above formula makes sense. Moreover,

$$\Big\|(H_0-\lambda)^{-1}V\Big\|_{p\to p} \le \Big\|(H_0-\lambda)^{-1}\Big\|_{p\to p}\|V\|_\infty \le c_1[(\mathrm{Re}\sqrt{\tau-\lambda})^{-2} + (\mathrm{Re}\sqrt{\lambda+\tau})^{-2}]\|V\|_\infty.$$

Hence if $(\mathrm{Re}\sqrt{\tau-\lambda})^2 \ge 4c_1\|V\|_\infty$, $(\mathrm{Re}\sqrt{\lambda+\tau})^2 \ge 4c_1\|V\|_\infty$, we have

$$\Big\|(H(\alpha)-\lambda)^{-1}\Big\|_{p\to p} \le 2\Big\|(H_0-\lambda)^{-1}\Big\|_{p\to p} \le c[(\mathrm{Re}\sqrt{\tau-\lambda})^{-2} + (\mathrm{Re}\sqrt{\lambda+\tau})^{-2}].$$

For $0<s<2$, $x \in H^2$, define

$$H(\alpha)^{s/2}x = \frac{1}{2\pi i}\int_\Gamma \lambda^{\frac{s}{2}-1}(\lambda - H(\alpha))^{-1}H(\alpha)x d\lambda. \tag{3.14}$$

where

$$\Gamma = \gamma_{1,+}\cup\gamma_{2,+}\cup\gamma_{1,-}\cup\gamma_{2,-},\ \gamma_{1,\pm}(t) = t+a\pm i\varepsilon t, t\ge 0,\ \gamma_{2,\pm}(t) = -t-a\pm i\varepsilon, t \ge 0.$$

See Figure 1 for the shape of Γ and the two connected branches of Γ are anticlockwise oriented. Here $a>0$ is some appropriate constant close to τ excluding discrete spectrum of $H(\alpha)$ in the interior of Γ, and ε is sufficiently small to ensure $\gamma_{1,\pm}(t)$ and $\gamma_{2,\pm}(t)$ lie in some single-value branch of $z^{s/2}$. It is a direct corollary that $H(\alpha)^{s/2}x = 0$ if x lies in the discrete part. The integration is well-defined for each $x \in H^2$, because of Lemma 3.2. Similarly, we define $H_0^{s/2}$.

For $\beta < 0$, define

$$[H(\alpha)]^\beta x = \frac{1}{2\pi i}\int_\Gamma \lambda^\beta(\lambda - H(\alpha))^{-1}x d\lambda.$$

For each $x \in L^2$, the integration is well-defined due to Lemma 3.2.



Figure 1: curve Γ

Lemma 3.3. *Denote $\mathcal{H}$ as the continuous spectral part of $H(\alpha)$ in L^2, then in $\mathcal{H}$*

$$H(\alpha)^{s/2} = ([H(\alpha)]^{-s/2})^{-1}.$$

Proof *Step one.* We claim $[H(\alpha)]^{-1}$ is exactly the inverse of $H(\alpha)$ in $\mathcal{H}$. It suffices to prove

$$\frac{1}{2\pi i}\int_\Gamma H(\alpha)(\lambda - H(\alpha))^{-1}\lambda^{-1}d\lambda = P_c(H(\alpha)). \tag{3.15}$$

For simplicity we denote $\xi_1(\alpha)$ ($\xi_2(\alpha)$) by z (y respectively). Then the discrete part of $H(\alpha)$ is spanned by {z,y} and $H(\alpha)z = 0$, $H(\alpha)y = iz$, it is easily seen

$$\int_\Gamma H(\alpha)(\lambda - H(\alpha))^{-1}\lambda^{-1}z d\lambda = 0$$
$$\int_\Gamma (\lambda - H(\alpha))^{-1}\lambda^{-1}H(\alpha)y d\lambda = \int_\Gamma i\lambda^{-2}z d\lambda = 0.$$

In other words, it vanishes in discrete part of $H(\alpha)$. Let

$$L = \frac{1}{2\pi i}\int_\Gamma H(\alpha)(\lambda - H(\alpha))^{-1}\lambda^{-1}d\lambda,$$

then in order to prove (3.15), we need to check the following two things:
(i)L is a projection;
(ii)$\langle Lv, \theta_3 w\rangle = 0$, for each $v \in L^2$ and w belonging to the discrete part of $H(\alpha)$.
First we show (ii) is correct. Indeed,

$$\left\langle \int_\Gamma H(\alpha)(\lambda - H(\alpha))^{-1}\lambda^{-1}d\lambda v, \theta_3 w\right\rangle$$
$$= \left\langle v, \int_\Gamma H(\alpha)^*(\bar{\lambda} - H(\alpha)^*)^{-1}\bar{\lambda}^{-1}\theta_3 w d\bar{\lambda}\right\rangle$$

$$
\begin{aligned}
&= \left\langle v, \theta_3 \int_{\Gamma} H(\alpha)(\lambda - H(\alpha))^{-1}\lambda^{-1} w d\lambda \right\rangle \\
&= 0.
\end{aligned}
$$

Second, we prove (i) is valid.



Figure 2: curve Γ and Γ'

Taking curve Γ' (see Figure 2)

$$\Gamma' = \gamma_{1,+} \cup \gamma_{2,+} \cup \gamma_{1,-} \cup \gamma_{2,-},\ \ \gamma_{1,\pm}(t) = t + a' \pm i\varepsilon t, t \geq 0,\ \ \gamma_{2,\pm}(t) = -t - a' \pm i\varepsilon t, t \geq 0,$$

where $a < a' < \tau$, we will show

$$\int_{\Gamma} H(\alpha)(\lambda - H(\alpha))^{-1}\lambda^{-1} d\lambda = \int_{\Gamma'} H(\alpha)(\lambda - H(\alpha))^{-1}\lambda^{-1} d\lambda.$$

Indeed, for $\beta < 0$, we can prove

$$\int_{\Gamma} \lambda^{\beta}(\lambda - H(\alpha))^{-1} d\tau = \int_{\Gamma'} \lambda^{\beta}(\lambda - H(\alpha))^{-1} d\lambda. \tag{3.16}$$

Denote

$$
\begin{aligned}
&\Gamma_{1,n} = \{-t - a + it\varepsilon : 0 \leq t \leq n\} \cup \{-t - a - it\varepsilon : 0 \leq t \leq n\} \\
&\Gamma'_{1,n} = \{-t - a' + it\varepsilon : 0 \leq t \leq n\} \cup \{-t - a' - it\varepsilon : 0 \leq t \leq n\} \\
&\gamma_{1,n} = \{-n + it : -\varepsilon(n + a) \leq t \leq -\varepsilon(n + a')\} \cup \{-n + it : \varepsilon(n + a') \leq t \leq \varepsilon(n + a)\},
\end{aligned}
$$

where $\gamma_{1,n}$ is oriented downward, $\Gamma_{1,n}$ and $\Gamma'_{1,n}$ are oriented anticlockwise. (see Figure 3) Then from the analyticity of resolvent,

$$\int_{\Gamma_{1,n}} \lambda^{\beta}(\lambda - H(\alpha))^{-1} x d\lambda - \int_{\Gamma'_{1,n}} \lambda^{\beta}(\lambda - H(\alpha))^{-1} x d\lambda + \int_{\gamma_{1,n}} \lambda^{\beta}(\lambda - H(\alpha))^{-1} x d\lambda = 0.$$



Figure 3: curve $\Gamma_{1,n}$, $\Gamma'_{1,n}$, and $\gamma'_{1,n}$

.

Since Γ and Γ' are symmetric respect to y axiom, it suffices to prove

$$\lim_{n\to\infty}\int_{\gamma_{1,n}} \lambda^{\beta}(\lambda - H(\alpha))^{-1}x d\lambda = 0. \tag{3.17}$$

From the decay of resolvent, for n sufficiently large, we have

$$\begin{aligned}
&\left\|\int_{\gamma_{1,n}} \lambda^{\beta}(\lambda - H(\alpha))^{-1}x d\lambda\right\|_p \\
&\leq \int_{-\varepsilon(n+a)}^{-\varepsilon(n+a')} \left\|(-n+it-H(\alpha))^{-1}x\right\|_p |-n+it|^{\alpha}dt + \int_{\varepsilon(n+a')}^{\varepsilon(n+a)} \left\|(-n+it-H(\alpha))^{-1}x\right\|_p |-n+it|^{\alpha}dt \\
&\leq \int_{-\varepsilon(n+a)}^{-\varepsilon(n+a')} \left(\mathrm{Re}\sqrt{\lambda \pm (it-n)}\right)^{-2} |-n+it|^{\alpha}dt\|x\|_p + \int_{\varepsilon(n+a')}^{\varepsilon(n+a)} \left(\mathrm{Re}\sqrt{\lambda \pm (it-n)}\right)^{-2} |-n+it|^{\alpha}dt\|x\|_p.
\end{aligned}$$

Since as $n \to \infty$, $\arg\left(\lambda \pm (it-n)\right) \sim \pm \arctan\varepsilon$, we find $\left(\mathrm{Re}\sqrt{\lambda \pm (it-n)}\right)^{-2} \sim c|n|^{-1}$, which immediately leads to (3.17). Then we have

$$\begin{aligned}
&\left(\int_{\Gamma} H(\alpha)(\lambda - H(\alpha))^{-1}\lambda^{-1}d\lambda\right)^2 \\
&= \int_{\Gamma}\int_{\Gamma'} H(\alpha)(\lambda - H(\alpha))^{-1}\lambda^{-1}H(\alpha)(\mu - H(\alpha))^{-1}\mu^{-1}d\lambda d\mu \\
&= \int_{\Gamma}\int_{\Gamma'} H(\alpha)^2\frac{\lambda^{-1}\mu^{-1}}{\mu-\lambda}\left[(\lambda - H(\alpha))^{-1} - (\mu - H(\alpha))^{-1}\right] d\lambda d\mu \\
&= 2\pi i\int_{\Gamma'} H(\alpha)^2\mu^{-2}(\mu - H(\alpha))^{-1}d\mu \\
&= -2\pi i\int_{\Gamma'} \mu^{-2}H(\alpha)d\mu + 2\pi i\int_{\Gamma'} \mu^{-1}(\mu - H(\alpha))^{-1}H(\alpha)d\mu \\
&= 2\pi i\int_{\Gamma'} \mu^{-1}(\mu - H(\alpha))^{-1}H(\alpha)d\mu
\end{aligned}$$

$$= 2\pi i \int_{\Gamma} H(\alpha)(\lambda - H(\alpha))^{-1}\lambda^{-1} d\lambda.$$

Thus L is a projection.

Step two. We prove for each $\gamma, \beta < 0$,

$$[H(\alpha)]^{\gamma+\beta} = [H(\alpha)]^{\gamma}[H(\alpha)]^{\beta}. \tag{3.18}$$

From (3.16), we deduce

$$\begin{aligned}
[H(\alpha)]^{\gamma}[H(\alpha)]^{\beta}x &= \frac{1}{(2\pi i)^2}\int_{\Gamma}\mu^{\gamma}(\tau - H(\alpha))^{-1}d\mu\int_{\Gamma'}\lambda^{\beta}(\lambda - H(\alpha))^{-1}xd\lambda \\
&= \frac{1}{(2\pi i)^2}\int_{\Gamma}\int_{\Gamma'}\lambda^{\beta}\mu^{\gamma}(\mu - H(\alpha))^{-1}(\lambda - H(\alpha))^{-1}xd\lambda d\mu \\
&= \frac{1}{(2\pi i)^2}\int_{\Gamma}(\mu - H(\alpha))^{-1}d\mu\int_{\Gamma'}\frac{\lambda^{\beta}\mu^{\gamma}}{\mu-\lambda}xd\lambda - \int_{\Gamma'}(\lambda - H(\alpha))^{-1}\int_{\Gamma}\frac{\lambda^{\beta}\mu^{\gamma}}{\mu-\lambda}xd\mu \\
&= \frac{1}{2\pi i}\int_{\Gamma'}(\lambda - H(\alpha))^{-1}\lambda^{\beta+\gamma}xd\lambda \\
&= [H(\alpha)]^{\gamma+\beta}x.
\end{aligned}$$

The following two steps are standard (see A. Pazy [12]), but for completeness, we give a sketch.

Step three. In this step, we define $[H(\alpha)]^{\gamma}$ for $\gamma \in \mathbb{R}$, and extend (3.18) to $\gamma, \beta \in \mathbb{R}$. From step one, we find $[H(\alpha)]^{-1}$ is one to one in $\mathcal{H}$. Therefore, for any integer $n \geq 1$, $[H(\alpha)]^{-n}$ is one to one. Suppose $[H(\alpha)]^{\gamma}x = 0$ and take $n \geq |\alpha|$, then from Step two, we have $[H(\alpha)]^{-n}x = [H(\alpha)]^{-n-\gamma}[H(\alpha)]^{\gamma}x = 0$. This implies $x = 0$. Thus $[H(\alpha)]^{\gamma}$ is one-to-one and we can define $[H(\alpha)]^{-gamma}$ as $([H(\alpha)]^{\gamma})^{-1}$. We claim $[H(\alpha)]^{\gamma}[H(\alpha)]^{\beta}x = [H(\alpha)]^{\gamma+\beta}x$ for $\gamma, \beta \in \mathbb{R}$ and $x \in D([H(\alpha)]^{\theta})$, where $\theta = \max(\gamma, \beta, \gamma+\beta)$. Indeed. for example, when $\gamma > 0$, $\beta < 0$, $\gamma + \beta > 0$, for $x \in D([H(\alpha)]^{\theta})$, in order to verify $[H(\alpha)]^{\gamma+\beta}x = [H(\alpha)]^{\gamma}[H(\alpha)]^{\beta}x$, it suffices to prove $[H(\alpha)]^{\beta}x = [H(\alpha)]^{-\gamma}[H(\alpha)]^{\gamma+\beta}x$. Let $y = [H(\alpha)]^{\gamma+\beta}x$, then it is equivalent to prove $[H(\alpha)]^{-\gamma}y = [H(\alpha)]^{\beta}[H(\alpha)]^{-\gamma-\beta}y$, which follows quickly from Step two.

Step four. In this step, we finish our proof. From Step three, $[H(\alpha)]^{s/2} = [H(\alpha)][H(\alpha)]^{s/2-1} = H(\alpha)[H(\alpha)]^{s/2-1}$, by the definition of $H(\alpha)^{s/2}$, $H(\alpha)^{s/2} = [H(\alpha)]^{s/2}$. Since $[H(\alpha)]^{s/2}[H(\alpha)]^{-s/2} = I$, we obtain our lemma.

Lemma 3.4. *For $2 \leq p \leq \infty$, it holds*

$$\left\| H(\alpha)^{s/2}f - H_0^{s/2}f \right\|_2 \leq C\left\| f \right\|_p, \tag{3.19}$$

Proof It is easy to see

$$(\lambda - H(\alpha))^{-1} - (\lambda - H_0)^{-1} = (\lambda - H(\alpha))^{-1}V(\lambda - H_0)^{-1}.$$

Therefore, from (3.14), we have

$$\begin{aligned}
H(\alpha)^{\frac{s}{2}} =& \int\limits_{\Gamma} H(\alpha)\lambda^{-1+\frac{s}{2}}(\lambda - H(\alpha))^{-1}d\lambda \\
=& \int\limits_{\Gamma} H(\alpha)\lambda^{-1+\frac{s}{2}}\left((\lambda - H(\alpha))^{-1} - (\lambda - H_0)^{-1}\right)d\lambda + H(\alpha)\int\limits_{\Gamma} \lambda^{-1+\frac{s}{2}}(\lambda - H_0)^{-1}d\lambda \\
=& \int\limits_{\Gamma} H(\alpha)\lambda^{-1+\frac{s}{2}}(\lambda - H(\alpha))^{-1}V(\lambda - H_0)^{-1}d\lambda + (H_0)^{\frac{s}{2}} + V\int\limits_{\Gamma} \lambda^{-1+\frac{s}{2}}(\lambda - H_0)^{-1}d\lambda \\
=& -\int\limits_{\Gamma} (\lambda - H(\alpha))\lambda^{-1+\frac{s}{2}}(\lambda - H(\alpha))^{-1}V(\lambda - H_0)^{-1}d\lambda + V\int\limits_{\Gamma} \lambda^{-1+\frac{s}{2}}(\lambda - H_0)^{-1}d\lambda \\
&+ \int\limits_{\Gamma} \lambda^{\frac{s}{2}}(\lambda - H(\alpha))^{-1}V(\lambda - H_0)^{-1}d\lambda + (H_0)^{\frac{s}{2}} \\
=& \int\limits_{\Gamma_1} \lambda^{\frac{s}{2}}(\lambda - H(\alpha))^{-1}V(\lambda - H_0)^{-1}d\lambda + (H_0)^{\frac{s}{2}}
\end{aligned}$$

Define

$$\begin{aligned}
\Gamma_1 &= \Gamma \cap \left\{ (\mathrm{Re}\sqrt{\tau - \lambda})^2 \geq 4c_1\|V\|_\infty, (\mathrm{Re}\sqrt{\lambda + \tau})^2 \geq 4c_1\|V\|_\infty \right\}, \\
\Gamma_2 &= \Gamma \cap \left\{ (\mathrm{Re}\sqrt{\tau - \lambda})^2 \leq 4c_1\|V\|_\infty, (\mathrm{Re}\sqrt{\lambda + \tau})^2 \geq 4c_1\|V\|_\infty \right\}, \\
\Gamma_3 &= \Gamma \cap \left\{ (\mathrm{Re}\sqrt{\tau - \lambda})^2 \geq 4c_1\|V\|_\infty, (\mathrm{Re}\sqrt{\lambda + \tau})^2 \leq 4c_1\|V\|_\infty \right\}, \\
\Gamma_4 &= \Gamma \cap \left\{ (\mathrm{Re}\sqrt{\tau - \lambda})^2 \leq 4c_1\|V\|_\infty, (\mathrm{Re}\sqrt{\lambda + \tau})^2 \leq 4c_1\|V\|_\infty \right\},
\end{aligned}$$

then from Lemma 3.2, Lemma 3.1, when $\frac{1}{q} + \frac{1}{p} = \frac{1}{2}$, we have

$$\begin{aligned}
\left\| \int\limits_{\Gamma_1} \lambda^{\frac{s}{2}}(\lambda - H(\alpha))^{-1}V(\lambda - H_0)^{-1}f d\lambda \right\|_2 &\leq \int\limits_{\Gamma_1} |\lambda|^{\frac{s}{2}} \left\|(\lambda - H(\alpha))^{-1}\right\|_{2\to 2}\|V\|_q \left\|(\lambda - H_0)^{-1}\right\|_{p\to p}\|f\|_p d\lambda \\
&\leq \int\limits_{\Gamma_1} |\lambda|^{\frac{s}{2}}[(\mathrm{Re}\sqrt{\tau - \lambda})^{-2} + (\mathrm{Re}\sqrt{\lambda + \tau})^{-2}]^2\|V\|_q\|f\|_p d\lambda.
\end{aligned} \tag{3.20}$$

The only singular point for (3.20) is infinity. But at infinity, the integrand behaves like $|\lambda|^{\frac{s}{2}-2}$, which is integrable. Noticing that $(\mathrm{Re}\sqrt{\tau - \lambda})^{-2} \sim |\lambda|$, $(\mathrm{Re}\sqrt{\lambda + \tau})^{-2} \sim |\lambda|$, as $\lambda \to \infty$ in Γ, the curve Γ_2, Γ_3, Γ_4 are actually bounded. Hence the remaining three terms can be estimated by

$$\int\limits_{\{\lambda:|\lambda|\leq R\}\cap\Gamma} |\lambda|^{\frac{s}{2}} \left\|(\lambda - H(\alpha))^{-1}\right\|_{2\to 2}\|V\|_q \left\|(\lambda - H_0)^{-1}\right\|_{p\to p}\|f\|_p d\lambda$$

$$\leq C\|V\|_q\|f\|_p.$$

Lemma 3.5. *For $1/2 < s < 2$, $2 \leq p \leq \infty$, it holds*

$$\|f\|_p + \|f\|_{H^s} \leq C\|H_0^{s/2} f\|_2$$

Proof From Fourier transformation, it is easy to see $H_0^{s/2}$ is roughly $(\tau-\Delta)^{s/2}$. Indeed, applying Fourier transformation, we have

$$\begin{aligned}
&\mathcal{F}\left[\int_\Gamma \lambda^{\frac{s}{2}-1}(\lambda - H_0)^{-1} H_0 f d\lambda\right](k) \\
&= \begin{pmatrix} \tau + |k|^2 & \\ & -\tau - |k|^2 \end{pmatrix} \int_\Gamma \lambda^{\frac{s}{2}-1} \begin{pmatrix} \left[\lambda - (\tau + |k|^2)\right]^{-1} & \\ & \left[\lambda + \tau + |k|^2\right]^{-1} \end{pmatrix} \mathcal{F}(f)(k) d\lambda \\
&= \begin{pmatrix} \left(\tau + |k|^2\right)^{s/2} & \\ & \left(-\tau - |k|^2\right)^{s/2} \end{pmatrix} \mathcal{F}(f)(k).
\end{aligned}$$

Standard Sobolev imbedding theorem yields $\|f\|_p \leq C\left\|H_0^{s/2} f\right\|_2$.

Lemma 3.6. *For f in the continuous spectral space of $H(\alpha)$, $1/2 < s < 2$, $2 \leq p \leq \infty$, we have*

$$\|f\|_p \leq C\|H(\alpha)^{s/2} f\|_2.$$

Proof It suffices to prove $\left\|[H(\alpha)]^{-s/2} f\right\|_p \leq C\|f\|_2$, due to Lemma 3.3. Similar arguments as Lemma 3.4 yield,

$$\begin{aligned}
\left\|[H(\alpha)]^{-s/2} f - [H_0]^{-s/2} f\right\|_p &\leq C\left\|\int_\Gamma \lambda^{-s/2+1}(\lambda - H_0)^{-1} V (\lambda - H(\alpha))^{-1} f d\lambda\right\|_p \\
&\leq C\int_\Gamma |\lambda|^{-s/2+1}\left\|(\lambda - H_0)^{-1}\right\|_{2\to p} \|V\|_\infty \left\|(\lambda - H(\alpha))^{-1}\right\|_{2\to 2} \|f\|_2 d\lambda.
\end{aligned}$$

Applying Lemma 3.2, Lemma 3.1 and Lemma 3.5, using similar arguments in Lemma 3.4, we obtain Lemma 3.6.

Lemma 3.6 and the definition of $|J_V|^s u$ immediately yield

Corollary 3.7. *If $U(-t)u$ belongs to the continuous part of $H(\alpha)$, $2 \leq p \leq \infty$ then*

$$\|u\|_p \leq t^{-s}\||J_V|^s u\|_{L^2},$$

where $|J_V|^s f = U(t)(t^2 H(\alpha))^{s/2} U(-t) f$.

Denote the projection to the continuous part of $H(\alpha(t))$ by P_1. Fix time $t_1 > h$, and denote

$\alpha(t_1)$ by α_1. Recall

$$H(\alpha_1) = \begin{pmatrix} -\Delta + \frac{\alpha_1^2}{4} & \\ & \Delta - \frac{\alpha_1^2}{4} \end{pmatrix} + \begin{pmatrix} V_1(\alpha_1) & V_2(\alpha_1) \\ V_2(\alpha_1) & -V_1(\alpha_1) \end{pmatrix}.$$

From Section 1, we know that the discrete spectral part of $H(\alpha_1)$ is spanned by $\{\xi_1(\alpha_1), \xi_2(\alpha_1)\}$. Moreover $\xi_1(\alpha_1) \to \xi_1(\alpha(t))$, $\xi_2(\alpha_1) \to \xi_2(\alpha(t))$, as $\alpha_1 \to \alpha(t)$. Define the projection to the continuous part of $H(\alpha_1)$ as P_2. Recall $\vec{g}$ in the decomposition of $u(x,t)$ in Section 2. For simplicity, we sometimes write $\vec{g}$ as f.

Lemma 3.8. *If $|\alpha_1 - \alpha(t)| \ll 1$, then for some constant C we have*

$$\| |J_V|^s g\|_2 \le C \|P_2 |J_V|^s g\|_2.$$

Proof Suppose $|J_V|^s g = P_2 |J_V|^s g + h_1 \xi_1(\alpha_1) + h_2 \xi_2(\alpha_1)$, then the orthogonal condition (2.7) gives

$$\begin{aligned} 0 &= \langle |J_V|^s g, U(t)\theta_3 \xi_i(\alpha(t))\rangle \\ &= \langle |J_V|^s g, U(t)\theta_3 \xi_i(\alpha_1)\rangle + \langle |J_V|^s g, U(t)\theta_3 [\xi_i(\alpha(t)) - \xi_i(\alpha_1)]\rangle \\ &= \langle |J_V|^s g, \theta_3 \xi_i(\alpha_1)\rangle + O\left(|\alpha_1 - \alpha|\right) \| |J_V|^s g\|_2 \end{aligned}$$

Therefore substituting the decomposition of $|J_V|^s f$ into the above formula yields

$$\begin{aligned} \|(h_1, h_2)\| &\le C\|P_2 |J_V|^s g\|_2 + O\left(|\alpha_1 - \alpha|\right) \| |J_V|^s g\|_2 \\ &\le C\|P_2 |J_V|^s g\|_2 + O\left(|\alpha_1 - \alpha|\right) \|(h_1, h_2)\| \end{aligned}$$

Thus if $|\alpha_1 - \alpha(t)| \ll 1$, we have

$$\|(h_1, h_2)\| \le C\|P_2 |J_V|^s g\|_2.$$

4 Nonlinear estimates

Define $\mathcal{M} = \sup_{\tau \le t_1} (\|f\|_m + \|f\|_n + \|f\|_\infty)\langle\tau\rangle^s$, since $s = \frac{7}{4}^+$, then from Lemma 2.2, we have

$$|\gamma'| + |\omega'| \le C W(\mathcal{M}) t^{-1-s} \mathcal{M}^2, \tag{4.21}$$

where $W(x)$ is some bounded function around zero. Thus

$$\sup_{\tau \le t_1} |\alpha(t) - \alpha_0| \le W(\mathcal{M}) \mathcal{M}^2,$$

$$\left|\alpha'\right| \leq W(\mathcal{M})t^{-1-s}\mathcal{M}^2, \tag{4.22}$$

$$|\alpha(t)-\alpha(t_1)| + |\Omega| \leq W(\mathcal{M})\mathcal{M}^2 t^{1-s}. \tag{4.23}$$

Furthermore if $\mathcal{M}$ is sufficiently small, then Corollary 3.7 and Lemma 3.8 yield

$$\mathcal{M} \leq C\|P_2|J_V|^s g\|_{L_t^\infty L_x^2}. \tag{4.24}$$

4.1 The equation for $|J_V|^s g$

Denote

$$\partial_t E_2 = \begin{pmatrix} \partial_t & \\ & \partial_t \end{pmatrix},$$

then direct calculations imply

Lemma 4.1.

$$\left[i\partial_t E_2 - \begin{pmatrix} -\Delta+\tau & \\ & \Delta-\tau \end{pmatrix}, \begin{pmatrix} M(t) & \\ & M(-t) \end{pmatrix}\right]$$
$$= \begin{pmatrix} M(t)\left(\frac{i}{2t}+\frac{ix\cdot\nabla}{t}\right) & \\ & M(-t)\left(\frac{i}{2t}+\frac{ix\cdot\nabla}{t}\right) \end{pmatrix}.$$

Lemma 4.2.

$$\left[i\partial_t E_2 - \begin{pmatrix} -\Delta+\tau & \\ & \Delta-\tau \end{pmatrix}, \begin{pmatrix} M(-t) & \\ & M(t) \end{pmatrix}\right]$$
$$= \begin{pmatrix} M(-t)\left(-\frac{i}{2t}-\frac{ix\cdot\nabla}{t}-\frac{x^2}{2t^2}\right) & \\ & M(t)\left(-\frac{i}{2t}-\frac{ix\cdot\nabla}{t}+\frac{x^2}{2t^2}\right) \end{pmatrix}.$$

For potential term, we have

Lemma 4.3.

$$[V, U(t)] = \begin{pmatrix} 0 & -V_2[M(-t)-M(t)] \\ V_2[M(-t)-M(t)] & 0 \end{pmatrix}.$$

For simplicity, we write $H(\alpha_1)$ as K.

Lemma 4.4. *If we define* $A = -U(t)\left(t^2K\right)^{s/2}[V, U(-t)] - [V, U(t)]\left(t^2K\right)^{s/2}U(-t)$, *then it holds that*

$$[i\partial_t E_2 - K, |J_V|^s] = \frac{is}{t}|J_V|^s + \frac{i}{t}U(t)\left[x\cdot\nabla, \left(t^2K\right)^{s/2}\right]U(-t) + A.$$

Proof Lemma 4.1 to Lemma 4.3 yield

$[i\partial_t E_2 - K, |J_V|^s]$

$$
\begin{aligned}
&= [i\partial_t E_2 - K, U(t)] \left(t^2K\right)^{s/2} U(-t) + U(t)\left[i\partial_t E_2 - K, \left(t^2K\right)^{s/2}U(-t)\right] \\
&= \frac{i}{2t}|J_V|^s + \begin{pmatrix} M(t)\,\frac{ix\cdot\nabla}{t} & \\ & M(-t)\,\frac{ix\cdot\nabla}{t} \end{pmatrix} \left(t^2K\right)^{s/2}U(-t) \\
&+ U(t)\left(t^2K\right)^{s/2}[i\partial_t E_2 - K, U(-t)] - [V, U(t)]\left(t^2K\right)^{s/2}U(-t) + U(t)[i\partial_t E_2 - K, (t^2K)^{s/2}]U(-t) \\
&= \begin{pmatrix} M(t)\,\frac{ix\cdot\nabla}{t} & \\ & M(-t)\,\frac{ix\cdot\nabla}{t} \end{pmatrix} \left(t^2K\right)^{s/2}U(-t) + A + \frac{is}{t}|J_V|^s + \frac{i}{2t}|J_V|^s \\
&+ U(t)\left(t^2K\right)^{s/2} \begin{pmatrix} M(-t)\,(-\frac{i}{2t} - \frac{ix\cdot\nabla}{t} - \frac{x^2}{2t^2}) & \\ & M(t)\,(-\frac{i}{2t} - \frac{ix\cdot\nabla}{t} + \frac{x^2}{2t^2}) \end{pmatrix} \\
&= \frac{is}{t}|J_V|^s + \begin{pmatrix} M(t)\,\frac{ix\cdot\nabla}{t} & \\ & M(-t)\,\frac{ix\cdot\nabla}{t} \end{pmatrix} \left(t^2K\right)^{s/2}U(-t) \\
&- U(t)\left(t^2K\right)^{s/2} \begin{pmatrix} M(-t)\,\frac{ix\cdot\nabla}{t} & \\ & M(t)\,\frac{ix\cdot\nabla}{t} \end{pmatrix} + U(t)\left(t^2K\right)^{s/2} \begin{pmatrix} M(-t)\,(-\frac{x^2}{2t^2}) & \\ & M(t)\,\frac{x^2}{2t^2} \end{pmatrix} + A \\
&= \frac{is}{t}|J_V|^s + \frac{i}{t}U(t)\left[x\cdot\nabla, \left(t^2K\right)^{s/2}U(-t)\right] + U(t)\left(t^2K\right)^{s/2} \begin{pmatrix} M(-t)\,(-\frac{x^2}{2t^2}) & \\ & M(t)\,\frac{x^2}{2t^2} \end{pmatrix} + A.
\end{aligned}
$$

Since

$$
\begin{aligned}
&\left[x\cdot\nabla, \left(t^2K\right)^{s/2}U(-t)\right] \\
&= \left[x\cdot\nabla, \left(t^2K\right)^{s/2}\right]U(-t) + \left(t^2K\right)^{s/2}[x\cdot\nabla, U(-t)] \\
&= \left[x\cdot\nabla, \left(t^2K\right)^{s/2}\right]U(-t) + \left(t^2K\right)^{s/2} \begin{pmatrix} M(-t)\,\frac{x^2}{2t^2} & \\ & -M(t)\,\frac{x^2}{2t^2} \end{pmatrix},
\end{aligned}
$$

our lemma follows.

If we denote

$$B(s) = s(K)^{s/2} + \left[x\cdot\nabla, (K)^{s/2}\right],$$

then

$$[i\partial_t E_2 - K, |J_V|^s] = it^{s-1}U(t)B(s)U(-t) + A + F. \tag{4.25}$$

For $B(s)$, we have

Lemma 4.5. *For $r = 1^+$,*

$$\|B(s)g\|_r \le C\|g\|_2,$$

Proof A small modification of the proof of Lemma 7.1 in [4] leads to

$$B(s) = c\int_\Gamma \tau^{s/2}(\tau - K)^{-1}V_3(\tau - K)^{-1}d\tau.$$

where $V_3 = 2V + x\frac{d}{dx}V$. Similar arguments as Lemma 3.4 yields our lemma. In fact, as presented in the proof of Lemma 3.4, we split Γ into four parts. Define

$$B_j g = \int_{\Gamma_j} \lambda^{s/2}(\lambda - K)^{-1} V_3 (\lambda - K)^{-1} g d\lambda.$$

Let $\frac{1}{r} = \frac{1}{2} + \frac{1}{q}$. For B_1, Lemma 3.2 and Hölder inequality give

$$\begin{aligned}
\|B_1 f\|_r &\leq \left\| \int_{\Gamma_1} \lambda^{s/2}(\lambda - K)^{-1} V_3 (\lambda - K)^{-1} g d\lambda \right\|_1 \\
&\leq \int_{\Gamma_1} \left\| (\lambda - K)^{-1} \right\|_{r \to r} \lambda^{s/2} \|V_3\|_q \left\| (\lambda - K)^{-1} \right\|_{2 \to 2} d\lambda \|g\|_2 \\
&\leq C \|g\|_2 \int_{\Gamma_1} \lambda^{s/2 - 2} d\lambda \\
&\leq C \|g\|_2.
\end{aligned}$$

Denote the resolvent set of K in L^2 by $\rho_{L^2}(K)$ and the resolvent set (spectrum) of K in $L^2 \cap L^1$ by $\rho_{L^2 \cap L^1}(K)$ ($\sigma_{L^2 \cap L^1}(K)$ respectively). The domain of K in $L^2 \cap L^r$ is taken as $W^{2,r} \cap W^{2,2}$. We claim $\rho_{L^2}(K) \subseteq \rho_{L^2 \cap L^r}(K)$. In fact, let $\vec{g} \in L^r \cap L^2$, $\lambda \in \rho_{L^2}(K)$, the definition of resolvent set gives $\vec{f} \in L^2$ for which it holds $\vec{g} = (K - \lambda)\vec{f}$ namely

$$\begin{cases} -\Delta f_1 + \tau_1 f_1 - \lambda f_1 + V_1(\alpha_1) f_1 - V_2(\alpha_1) f_2 = g_1 \\ \Delta f_2 - \tau_1 f_2 - \lambda f_2 + V_2(\alpha_1) f_1 - V_1(\alpha_1) f_2 = g_2 \end{cases} \tag{4.26}$$

Hölder inequality implies $g_1 - V_1(\alpha_1) f_1 + V_2(\alpha_1) f_2 \in L^r$. Therefore we have

$$-\Delta f_1 + (\tau_1 - \lambda) f_1 \in L^r.$$

Since $\tau_1 - \lambda \notin (-\infty, 0]$, $-\Delta f_1 + (\tau_1 - \lambda)$ is invertible in L^r, which can be proved by the integral expression of the resolvent of Δ. Hence $f_1 \in L^r$, again by the first equation in (4.26) we have $\Delta f_1 \in L^r$, thus $f_1 \in W^{2,r}$. The same arguments show $f_2 \in W^{2,r}$, by which we have proved our claim. In other words, we have proved $\sigma_{L^2 \cap L^r}(K) \subseteq \sigma_{L^2}(K)$, which indicates for $\lambda \in \Gamma_2 \cup \Gamma_3 \cup \Gamma_4$,

$$\left\| (\lambda - K)^{-1} \right\|_{L^r \cap L^2 \to L^r \cap L^2} \leq C.$$

Hence by Hölder inequality and the boundedness of Γ_j where $j \in \{2, 3, 4\}$, we obtain

$$\begin{aligned}
\|B_j g\|_1 &\leq \left\| \int_{\Gamma_j} \lambda^{s/2}(\lambda - K)^{-1} V_3 (\lambda - K)^{-1} g d\lambda \right\|_1 \\
&\leq \int_{\Gamma_j} \lambda^{s/2} \left\| (\lambda - K)^{-1} \right\|_{L^r \cap L^2 \to L^r \cap L^2} \left(\|V_3\|_q + \|V_3\|_\infty \right) \left\| (\lambda - K)^{-1} \right\|_{2 \to 2} d\lambda \|g\|_2 \\
&\leq C \|g\|_2.
\end{aligned}$$

Lemma 4.6. *Let* $r = 1^+$*, then*

$$\left\| K^{s/2} g - H_0^{s/2} g \right\|_r \leq C \|g\|_2.$$

Proof From the proof of Lemma 3.4,

$$\left\| K^{s/2} g - H_0^{s/2} g \right\|_r = \left\| \int_\Gamma \lambda^{s/2} (\lambda - K)^{-1} V (\lambda - H_0)^{-1} g \right\|_r.$$

The same arguments in the proof of Lemma 4.5 yield our Lemma.

Lemma 4.7. *Let* $r = 1^+$*,* (p, r) *is an admissible pair. Then*

$$\|Ag\|_{L_t^p L_x^r} \leq C(h^{-1}) \|P_2 |J_V|^s g\|_{L_t^\infty L_x^2}.$$

Proof Let $\frac{1}{r} = \frac{1}{2} + \frac{1}{q}$. Lemma 4.6, fractional Leibnitz formula, Lemma 3.4, Corollary 3.7 and Lemma 3.8 imply

$$\begin{aligned}
\|Ag\|_r &\leq \left\| t^s H_0^{s/2} [U(t) \breve{V}(t) U(-t) g] \right\|_r + \left\| \breve{V}(t) |J_V|^s g \right\|_r + t^s \left\| \breve{V}(t) g \right\|_2 \\
&\leq \left\| t^s U(t) \breve{V}(t) U(-t) g \right\|_{W^{s,r}} + \left\| \widehat{V}(t) |x|^2 \right\|_q \| |J_V|^s g\|_2 t^{-1} + t^{s-1} \|g\|_2 \left\| \widehat{V}(t) |x|^2 \right\|_\infty \\
&\leq \left\| t^s U(t) \breve{V}(t) \right\|_{W^{s,2}} \|g\|_q + \|t^s U(-t) g\|_{W^{s,2}} \left\| \breve{V}(t) \right\|_q + C \| |J_V|^s g\|_2 t^{-1} + C t^{s-1} \|g\|_2 \\
&\leq C t^{s-1} \|g\|_q + C t^{-1} \| |J_V|^s g\|_2 + C t^{s-1} \|g\|_2 \\
&\leq C \| |J_V|^s g\|_2 t^{-1} \leq C \|P_2 |J_V|^s g\|_{L_t^\infty L_x^2} t^{-1},
\end{aligned}$$

where

$$\breve{V}(t) = [M(t) - M(-t)] \begin{pmatrix} 0 & -V_2 \\ V_2 & 0 \end{pmatrix}, \ \widehat{V}(t) = \begin{pmatrix} 0 & -V_2 \\ V_2 & 0 \end{pmatrix}.$$

Lemma 4.8. *Let* $r = 1^+$*,* (p, r) *is an admissible pair. Define* $it^{s-1} U(t) B(s) U(-t) g = \bar{B}(s) g$*, then*

$$\left\| \bar{B}(s) g \right\|_{L_t^p L_x^r} \leq C(h^{-1}) \|P_2 |J_V|^s g\|_{L_t^\infty L_x^2}.$$

Proof Thanks to Lemma 4.5, Corollary 3.7 and Lemma 3.8, it is easy to see

$$\begin{aligned}
\left\| it^{s-1} P_2 U(t) B(s) U(-t) g \right\|_{L_t^p L_x^r} &\leq \left\| it^{s-1} g \right\|_{L_t^p L_x^2} \leq \left\| t^{s-1} t^{-s} \right\|_{L_t^r} \| |J_V|^s g\|_{L_t^\infty L_x^2} \\
&\leq C(h^{-1}) \|P_2 |J_V|^s h(t)\|_{L_t^\infty L_x^2}.
\end{aligned}$$

Lemma 4.9. *Let* $r = 1^+$*,* (p, r) *is an admissible pair. Then*

$$\left\| |J_V|^s \vec{D}_0 \right\|_{L_t^p L_x^r} \leq C \left\| P_2 |J_V|^s g \right\|_{L_t^\infty L_x^2}^2.$$

$$\begin{aligned}
\left\| |J_V|^s \vec{D}_4 \right\|_{L_t^p L_x^r} &\le C \left\| P_2 |J_V|^s g \right\|^2_{L_t^\infty L_x^2} + \left\| P_2 |J_V|^s g \right\|^{m-1}_{L_t^\infty L_x^2} + \left\| P_2 |J_V|^s g \right\|^{m-2}_{L_t^\infty L_x^2} . \\
\left\| |J_V|^s \vec{D}_j \right\|_{L_t^p L_x^r} &\le C \| P_2 |J_V|^s \vec{g} \|^3_{L_t^\infty L_x^2},
\end{aligned}$$

where $j = 1, 2, 3$.

Proof From (4.21),

$$\left\| |J_V|^s \vec{D}_0 \right\|_{L_t^p L_x^r} \le C \left\| |\gamma'|\, t^s + |\omega'|\, t^s \right\|_{L_t^p} \le C\mathcal{M}^2 \|t^{-1}\|_{L_t^p} \le C\mathcal{M}^2.$$

We can write D_4 as $D_4 = e^{-i\Omega} \int_0^1 [A_1 + A_2 + A_3 + A_4 + A_5](1-\tau)d\tau$, where

$$\begin{aligned}
A_1 &= \tilde{g}^2 \left(\overline{\varphi + \tau\tilde{g}}\right)^2 (\varphi + \tau\tilde{g}) F''(|\varphi + \tau\tilde{g}|^2), \\
A_2 &= 2|\tilde{g}|^2 (\varphi + \tau\tilde{g})^2 \overline{\varphi + \tau\tilde{g}} F''(|\varphi + \tau\tilde{g}|^2), \\
A_3 &= (\bar{\tilde{g}})^2 (\varphi + \tau\tilde{g})^3 F''(|\varphi + \tau\tilde{g}|^2), \\
A_4 &= 2\tilde{g}^2 \overline{\varphi + \tau\tilde{g}} F'(|\varphi + \tau\tilde{g}|^2), \\
A_5 &= 4|\tilde{g}|^2 (\varphi + \tau\tilde{g}) F'(|\varphi + \tau\tilde{g}|^2), \\
\tilde{g} &= g e^{i\Omega}.
\end{aligned}$$

We take A_3 as an example to illustrate how to bound them. Define $\frac{1}{r} = \frac{1}{2} + \frac{1}{q}$, $\{\eta\} \equiv (\eta, \bar{\eta})^t$. Then from Lemma 4.6, fractional Leibnitz formula, Lemma 3.5, Proposition 1.1, we obtain

$$\begin{aligned}
&\left\| |J_V|^s \left\{ (\bar{\tilde{g}})^2 (\varphi + \tau\tilde{g})^3 F''(|\varphi + \tau\tilde{g}|^2) \right\} \right\|_r \\
&\le t^s \left\| U(-t) \left\{ (\bar{\tilde{g}})^2 (\varphi + \tau\tilde{g})^3 F''(|\varphi + \tau\tilde{g}|^2) \right\} \right\|_{W^{s,r}} + Ct^s \left\| \left\{ (\bar{\tilde{g}})^2 (\varphi + \tau\tilde{g})^3 F''(|\varphi + \tau\tilde{g}|^2) \right\} \right\|_2 \\
&\le t^s \left\| M(-3t)(\varphi + \tau\tilde{g})^3 F''(|\varphi + \tau\tilde{g}|^2)(M(t)\bar{g})(M(t)\bar{g}) \right\|_{W^{s,r}} + Ct^s \|g\|^2_\infty \\
&\le t^s \left\| M(-3t)(\varphi + \tau\tilde{g})^3 F''(|\varphi + \tau\tilde{g}|^2) \right\|_{H^2} \left\| g^2 \right\|_q + t^s \left\| M(-3t)(\varphi + \tau\tilde{g})^3 F''(|\varphi + \tau\tilde{g}|^2) \right\|_q \left\| (M(t)\bar{g})^2 \right\|_{H^s} \\
&\quad + Ct^s \|g\|^2_\infty \\
&\le t^s \left\| M(-3t)(\varphi + \tau\tilde{g})^3 F''(|\varphi + \tau\tilde{g}|^2) \right\|_{H^2} \|g\|_\infty \|g\|_q + Ct^s \|M(t)\bar{g}\|_{H^s} \|g\|_\infty \\
&\quad + Ct^s \|g\|^2_\infty \\
&\le t^s \left\| M(-3t)(\varphi + \tau\tilde{g})^3 F''(|\varphi + \tau\tilde{g}|^2) \right\|_{H^2} \|g\|_\infty \|g\|_q + C \| |J_V|^s g \|_2 \|g\|_\infty + Ct^s \|g\|^2_\infty \\
&\le t^{s-2} \left\| |x|^2 (\varphi + \tau\tilde{g})^3 F''(|\varphi + \tau\tilde{g}|^2) \right\|_2 \|g\|_\infty \|g\|_q + t^{s-1} \left\| |x| \left((\varphi + \tau\tilde{g})^3 F''(|\varphi + \tau\tilde{g}|^2) \right)' \right\|_2 \|g\|_\infty \|g\|_q \\
&\quad + t^s \left\| \left((\varphi + \tau\tilde{g})^3 F''(|\varphi + \tau\tilde{g}|^2) \right)'' \right\|_2 \|g\|_\infty \|g\|_q + C \| |J_V|^s \vec{g} \|_2 \|g\|_\infty + Ct^s \|g\|^2_\infty \\
&\le t^{s-2} \left(\|g\|_\infty \|g\|_q + \left\| |x|^2 g \right\|_2 \|g\|_q \left(\|g\|^{m-2}_\infty + \|g\|^{n-2}_\infty \right) \right) + t^{s-1} \left(\|g\|_\infty \|g\|_q + \| |x|\, g \|_2 \|g\|_q \left(\|g\|^{m-3}_\infty + \|g\|^{n-3}_\infty \right) \right) \\
&\quad + C \| |J_V|^s \vec{g} \|_2 \|g\|_\infty + Ct^s \|g\|^2_\infty + Ct^s \|g\|_\infty \|g\|_q
\end{aligned}$$

$$
\begin{aligned}
&\lesssim t^{-s}\left\||J_V|^s\vec{g}\right\|_2^2 + t^{s+2-s(m-1)}\left\||J_V|^s\vec{g}\right\|_2^{m-1} + t^{s-(m-2)s}\left\||J_V|^s\vec{g}\right\|_2^{m-2} \\
&\quad + t^{s+2-s(n-1)}\left\||J_V|^s\vec{g}\right\|_2^{n-1} + t^{s-(n-2)s}\left\||J_V|^s\vec{g}\right\|_2^{n-2}.
\end{aligned}
$$

Since $s = \frac{7}{4}^{+}$, $m, n > \frac{26}{7}$, we get the estimates of D_4 in Lemma 4.9. Since the proofs of D_1, D_2, D_3 are almost the same, we only give the proof of D_3. Lemma 4.6, fractional Leibnitz formula and Lemma 3.5 give

$$
\begin{aligned}
&\left\||J_V|^s\vec{D}_3\right\|_r \\
&\leq t^s\left\|K^{s/2}U(-t)\vec{D}_3\right\|_r \\
&\leq Ct^s\left\|\int_{\alpha_1}^{\alpha(t)} U(-t)\left\{[F''(\varphi^2(\tau))\varphi^3(\tau)\varphi_\alpha(\tau) + F'(\varphi^2(\tau))\varphi(\tau)\varphi_\alpha(\tau)]\bar{g}\right\}d\tau\right\|_{W^{s,r}} \\
&\quad + Ct^s\left\|\int_{\alpha_1}^{\alpha(t)} U(-t)\left\{[F''(\varphi^2(\tau))\varphi^3(\tau)\varphi_\alpha(\tau) + F'(\varphi^2(\tau))\varphi(\tau)\varphi_\alpha(\tau)]\bar{g}\right\}d\tau\right\|_2 \\
&\leq Ct^s\int_{\alpha_1}^{\alpha(t)}\left\|M(-2t)[F''(\varphi^2(\tau))\varphi^3(\tau)\varphi_\alpha(\tau) + F'(\varphi^2(\tau))\varphi(\tau)\varphi_\alpha(\tau)]M(t)\bar{g}\right\|_{W^{s,r}}d\tau + \\
&\quad + Ct^s\left|\alpha(t)-\alpha_1\right|\|g\|_2 \\
&\leq Ct^s\int_{\alpha_1}^{\alpha(t)}\left\|M(-2t)[F''(\varphi^2(\tau))\varphi^3(\tau)\varphi_\alpha(\tau) + F'(\varphi^2(\tau))\varphi(\tau)\varphi_\alpha(\tau)]\right\|_{W^{s,2}}\|g\|_q d\tau \\
&\quad + Ct^s\int_{\alpha_1}^{\alpha(t)}\left\|M(-2t)[F''(\varphi^2(\tau))\varphi^3(\tau)\varphi_\alpha(\tau) + F'(\varphi^2(\tau))\varphi(\tau)\varphi_\alpha(\tau)]\right\|_q\|M(t)\bar{g}\|_{W^{s,2}}d\tau \\
&\quad + Ct^s\left|\alpha(t)-\alpha_1\right|\|g\|_2 \\
&\leq Ct^s\left|\alpha(t)-\alpha_1\right|\|g\|_q + C\left|\alpha(t)-\alpha_1\right|\left\||J_V|^s\vec{g}\right\|_2 + Ct^s\left|\alpha(t)-\alpha_1\right|\|g\|_2 \\
&\leq C\left|\alpha(t)-\alpha_1\right|\left\||J_V|^s\vec{g}\right\|_2 \\
&\leq Ct^{-s+1}\left\||J_V|^s\vec{g}\right\|_{L_t^\infty L_x^2}^3,
\end{aligned}
$$

where we have (4.23) in the last inequality.

4.2 Proof of main Theorem

Applying $|J_V|^s$ and P_2 to equation (2.13), then by commutator relations (4.25), we obtain

$$
i\partial_t P_2|J_V|^s\vec{g} - KP_2|J_V|^s\vec{g} - P_2|J_V|^s\vec{D} = P_2\left(\bar{B}(s)\vec{g} + A\vec{g}\right).
$$

From Lemma 4.7, Lemma 4.8, Lemma 4.9, and Strichartz estimates (see J. Krieger, W. Schlag [10]), we have

$$
\|P_2|J_V|^s g\|_{L_t^\infty L_x^2} \lesssim \left\||J_V|^s\vec{D}\right\|_{L_t^p L_x^r} + \left\|\bar{B}(s)g + Ag\right\|_{L_t^p L_x^r} + \|g_h\|_\Sigma
$$

$$\lesssim \|P_2|J_V|^s g\|^3_{L_t^\infty L_x^2} + \|P_2|J_V|^s g\|^2_{L_t^\infty L_x^2} + \|P_2|J_V|^s g\|^{m-1}_{L_t^\infty L_x^2} + \|P_2|J_V|^s g\|^{m-2}_{L_t^\infty L_x^2}$$
$$+ \|P_2|J_V|^s g\|^{n-1}_{L_t^\infty L_x^2} + \|P_2|J_V|^s g\|^{n-2}_{L_t^\infty L_x^2} + C(h^{-1})\|P_2|J_V|^s g\|_{L_t^\infty L^2} + C\|g_h\|_\Sigma.$$

If h is large enough, $\|g_h\|_\Sigma$ is sufficiently small, from standard continuity method, we obtain

$$\mathcal{M} \le \|P_2\,|J_V|\,g\|_{L_t^\infty L_x^2} < C.$$

Thus (4.21) implies, γ, ω have limits $\gamma_\infty, \omega_\infty$. Consequently, we can introduce the limit trajectory $\sigma_+(t)$:

$$\beta_+(t) = \omega_+ t + \gamma_+, \omega_+ = \omega_\infty, \gamma_+ = \int_0^\infty (\omega(\tau) - \omega_\infty) d\tau.$$

Obviously, $\sigma(t) - \sigma_+(t) = O(t^{-s+1})$ as $t \to \infty$. Hence, the limit soliton $w(x;\sigma_+(t))$ aries, and

$$w(x;\sigma(t)) - w(x;\sigma_+(t)) = O(t^{-s+1})$$

in the space of $L^2 \bigcap L^\infty$. Introduce transformation $\chi = e^{i\Phi_\infty} g(x,t)$, $\Phi_\infty = -\beta_+(t)$. Repeat the construction in section 2 to section 4, with the operator K, P_2 replaced by K_+, $P_c(K_+)$, where

$$K_+ = \begin{pmatrix} -\Delta - \omega_+ & \\ & \Delta + \omega_+ \end{pmatrix} + \begin{pmatrix} V_1(\alpha_+) & -V_2(\alpha_+) \\ V_2(\alpha_+) & -V_1(\alpha_+) \end{pmatrix}.$$

Then we can also prove $\|\chi\|_\infty \le Ct^{-s}$, therefore, we have obtained

$$u = w(x;\sigma_+(t)) + \chi + O(t^{-s+1}).$$

Combined with the estimate of χ, Theorem 1.1 follows.

5 Appendix A

First we prove $\|u|x|\|_2 \le C(t+1)$. ux satisfies

$$i\partial_t(xu) = -(\Delta u)\,x - F(|u|^2)ux, \tag{5.27}$$

Multiplying (5.27) by $\bar{u}x$, then taking the imaginary part, we have

$$\frac{1}{2}\frac{d}{dt}\||x|\,u\|_2^2 = 2\mathrm{Im}\int_{\mathbb{R}} (\nabla u)\, x\bar{u}.$$

Namely,

$$\frac{d}{dt}\||x|\,u\|_2^2 \le C\||x|\,u\|_2.$$

Gronwall's inequality yields $\|u|x|\|_2 \le C(t+1)$. Second, we prove $\|u'|x|\|_2 \le C(1+t^3)$. The equation for $u'x$ is

$$i\partial_t(xu') = -\left(\Delta u'\right)x - F'(|u|^2)(u'\bar{u} + \bar{u}'u)ux - F(|u|^2)u'x.$$

Multiplying the above formula with $\bar{u}'x$, taking the imaginary part, we obtain

$$\frac{d}{dt}\left\||x|\,u'\right\|_2^2 \lesssim \left\|u''\right\|_2\left\||x|\,u'\right\|_2 + \left\|u'\right\|_\infty^2\left(\|u\|_\infty^{m-3} + \|u\|_\infty^{n-3}\right)\||x|\,u\|_2^2.$$

Since $\|u\|_{H^2}$ is bounded, from Sobolev imbedding theorem, it follows,

$$\frac{d}{dt}\left\||x|\,u'\right\|_2^2 \le C\left\||x|\,u'\right\|_2 + Ct^2 + C.$$

Therefore,

$$\frac{d}{dt}\left(\left\||x|\,u'\right\|_2^2 + 1\right) \le C(t^2+2)\left(\left\||x|\,u'\right\|_2^2 + 1\right)^{1/2}.$$

Gronwall's inequality implies $\||x|\,u'\|_2 \le C(t^3+1)$. Third, we prove $\|u|x|^2\|_2 \le C(t^4+1)$. Similar arguments as the first step, we can show

$$\frac{d}{dt}\left\||x|^2u\right\|_2^2 \le C\left|\int_{\mathbb{R}} u'\bar{u}x^3\right| \le C\left\||x|^2u\right\|_2\||x|\,u'\|_2 \le C(t^3+1)\left\||x|^2u\right\|_2.$$

Again by Gronwall inequality, we get the desired result.

References

[1] M. Becenu, A critical center-stable manifold for Schrödinger's equation in three dimensions. Comm. Pure and App. Math. 65 (2012), 431-507.

[2] H. Berestycki, P.L. Lions, Nonlinear scalar field equations. I. Existence of a ground state. Arch. Ration. Mech. Anal. 82 (1983), 313-345.

[3] V.S. Buslaev, G.S. Perelman, Scattering for the nonlinear Schrödinger equation: states close to a soliton, St. Petersburg Math. J. 4 (1993), 1111-1142.

[4] S. Cuccagna, V. Georgiev, N. Visciglia. Decay and Scattering of small solutions of Pure Power NLS in R with $p > 3$ and with a potential. Comm. Pure Appl. Math. 6(2013), 957-980.

[5] S. Cuccagna, Stabilization of solutions to nonlinear Schrödinger equations, Comm. Pure App. Math. 54 (2001), 1110-1145.

[6] S. Cuccagna, An invariant set in enregy space for supercritical NLS in 1D. J. Math. Anal. Appl. 352 (2009), 634-644.

[7] S. Cuccagna, T. Mizumachi, On Asymptotic Stability in Energy Space of Ground States for Nonlinear Schrödinger Equations, Communications in Mathematical Physics, 284 (2008), pp 51-77.

[8] S. Gustafson, K. Nakanishi, T.P. Tsai, Asymptotic Stability and Completeness in the Energy Space for Nonlinear Schrödinger Equations with Small Solitary Waves, Int. Math. Res. Notices 66 (2004), 3559-3584.

[9] E. Kirr, Ö. Mizrak. Asymptotic stability of ground states in 3D nonlinear Schrödinger equation including subcritical cases, J. Func. Anal. 257 (2009) 3691-3747.

[10] J. Krieger, W. Schlag, Stable manifolds for all monic supercritical focusing nonlinear Schrödinger equations in one dimension, J. Amer. Math. Soc. 19 (2006) 815-920.

[11] E. Kirr, A. Zarnescu, On the Asymptotic Stability of Bound States in 2D Cubic Schrödinger Equation, Communications in Mathematical Physics, 272 (2007), pp 443-468.

[12] A. Pazy, Semigroups of Linear Operators and Applications to Partial Differential Equations, Springer-Verlag, 1983.

[13] G.S. Perelman, Asymptotic stability of solitons for nonlinear Schrödinger equations. Comm. in PDE, 29, 1051-1095 (2004)

[14] I. Rodnianski, W. Schlag, A. Soffer, Asymptotic stability of N-soliton states of NLS. http://arxiv.org/abs/math.AP/0309114, 2003.

[15] I. M. Sigal, Nonlinear wave and Schrödinger equations, I. Instability of periodic and quasiperiodic solutions. Commun. Math. Phys. 153, 297-320 (1993).

[16] Schlag W., Stable manifolds for an orbitally unstable nonlinear Schrödinger equation. Ann. of Math. 169 (2009), 139-227.

[17] A. Soffer, M. I. Weinstein. Multichannel nonlinear scattering theory for nonintegrable equations, Commun. Math. Phys. 342, 312-327 (1989).

[18] T. Mizumachi, Asymptotic stability of small solitary waves to 1D nonlinear Schrödinger equations with potential, J. Math. Kyoto Univ., Volume 48 (2008), 471-497.

[19] Weinstein M. I., Modulational stability of ground states of nonlinear Schrödinger equations. SIAM. J. Math. Anal. 16 (1985), 472-491.

[20] Weinstein M. I., Lyapunov stability of ground states of nonlinear dispersive evolution equations. Comm. Pure Appl. Math. 39 (1986), 51-67.